




European Research Council  
Established by the European Commission



# Indirect measurements of neutron-induced reaction cross sections at storage rings

**B. Jurado**<sup>1</sup>, M. Sguazzin<sup>1</sup>, J. Pibernat<sup>1</sup>, J. A. Swartz<sup>1</sup>, M. Grieser<sup>2</sup>, J. Glorius<sup>3</sup>, Y. A. Litvinov<sup>3</sup>, R. Reifarh<sup>4</sup>, K. Blaum<sup>2</sup>, P. Alfaut<sup>1</sup>, P. Ascher<sup>1</sup>, D. Atanasov<sup>1</sup>, L. Audouin<sup>5</sup>, C. Berthelot<sup>1</sup>, B. Blank<sup>1</sup>, B. Bruckner<sup>4</sup>, S. Dellmann<sup>4</sup>, I. Dillmann<sup>6</sup>, C. Domingo-Pardo<sup>7</sup>, M. Dupuis<sup>8</sup>, P. Erbacher<sup>4</sup>, M. Flayol<sup>1</sup>, O. Forstner<sup>3</sup>, D. Freire-Fernandez<sup>2</sup>, M. Gerbaux<sup>1</sup>, J. Giovinozzo<sup>1</sup>, S. Grevy<sup>1</sup>, C. Griffin<sup>6</sup>, A. Gumberidze<sup>3</sup>, S. Heil<sup>4</sup>, A. Heinz<sup>9</sup>, W. Korten<sup>8</sup>, D. Kurtulgil<sup>4</sup>, G. Leckenby<sup>6</sup>, S. Litvinov<sup>3</sup>, B. Lorentz<sup>3</sup>, V. Meot<sup>8</sup>, J. Michaud<sup>1</sup>, S. Perard<sup>1</sup>, U. Popp<sup>3</sup>, M. Roche<sup>1</sup>, M.S. Sanjari<sup>3</sup>, R.S. Sidhu<sup>10</sup>, U. Spillmann<sup>3</sup>, M. Steck<sup>3</sup>, Th. Stöhlker<sup>3</sup>, B. Thomas<sup>1</sup>, L. Thulliez<sup>8</sup>, M. Versteegen<sup>1</sup>

*1- LP2I (ex-CENBG), Bordeaux, France*

*2- MPIK, Heidelberg, Germany*

*3-GSI, Darmstadt, Germany*

*4-University of Frankfurt, Germany*

*5-IJCLAB, Orsay, France*

*6-Triumf, Vancouver, Canada*

*7-IFIC, Valencia, Spain*


*8-CEA, France*

*9-University of Chalmers, Sweden*

*10-University of Edinburgh, UK*


# Introduction:

## Neutron-induced reactions at energies below few MeV:



# Motivation:


## Need for neutron-induced reaction cross sections of radioactive nuclei



Essential for astrophysics, energy production and medicine!

# Need for neutron-induced reaction cross sections of short-lived nuclei


Synthesis of heavy elements via the s and r-processes



Fission sets the end point of the r-process path.  
Fission recycling.


- Very difficult or even impossible to measure with standard techniques because of the radioactivity of the targets.
- Complicated to calculate due to the difficulty to describe the de-excitation process. Calculations can be wrong by several orders of magnitude!

# Surrogate-reaction method



Decay probabilities as a function of excitation energy are precious observables to constrain model parameters (fission barriers, level densities...) and provide much more accurate predictions for neutron-induced cross-sections of nuclei far from stability.

# Benchmark:



First simultaneous  
measurement of  $P_f$  and  $P_\gamma$ !


Stringent test of  
experimental method!

Only way to access the  
fission threshold of fissile  
nuclei!


# Determination of model parameters

$$P_{s,decay}(E^*) = \sum_{J^\pi} F_s(E^*, J^\pi) G_{decay}(E^*, J^\pi)$$


Calculation of  $F_s$  populated in  
 $4\text{He} + 240\text{Pu} \rightarrow 4\text{He}' + 240\text{Pu}$   
 Marc Dupuis (CEA-DAM)



Statistical-model for de-excitation process (TALYS):



We need parameters for:  
 Fission barriers  
 Level densities  
 $\gamma$ -ray strength functions!




Knowing both  $P_f$  and  $P_\gamma$  below  $S_n$  has allowed us to determine these parameters precisely!

$$B_f = 5.98 \pm 0.02 \text{ MeV}$$


(typical uncertainty for  $B_f$  is 200 keV!)

# First simultaneous determination of neutron-induced fission and capture cross sections $n+^{239}\text{Pu}\rightarrow^{240}\text{Pu}^*$

## Fission




## Radiative capture





# Measurement of fission and gamma-emission probabilities in direct kinematics




## Limits:

- Unavailability of targets (radioactive samples)
- Target contaminants and target support
- $P_\gamma$ : rather low detection efficiency
- $P_n$ : measurement of low-energy neutrons and neutron efficiency

## Advantages of Inverse kinematics:

- Access to very short-lived nuclei
- Detection of heavy residues




## BUT!

- Required  $E^*$  resolution  $\sim$  few 100 keV,  
 $E^* = f(E_{\text{beam}}, E_{\text{target\_like}}, \theta)$
- Target contaminants and target windows have to be avoided

**STORAGE RINGS!**

# Advantages of heavy-ion storage rings


The ESR at GSI/FAIR





- Beam cooling → Excellent energy and position resolution of the beam, maintained after each passage through the target, negligible, E-loss & straggling effects
- Use of ultra-thin in-ring gas-jet targets  $\sim 10^{13}/\text{cm}^2$ . Effective target thickness increased by  $\sim 10^6$  due to revolution frequency (at 10 A MeV)
- High-quality, pure, fully-stripped beams and pure, ultra-thin, windowless targets → **unique!**

**Challenge: Detectors in Ultra-High Vacuum ( $10^{-11}$ - $10^{-12}$  mbar)!**

# First proof of principle experiment at the ESR, $^{208}\text{Pb}(p,p')$ , 20-27 June 2022




# Preliminary results, excitation energy resolution




$\Delta E^* \approx 600$  keV, dominated by the angular uncertainty due to target radius of 2.5 mm.  
With target radius 0.5 -1.5 mm  $\rightarrow \Delta E^* = 200-300$  keV!

# Preliminary results, detection of beam-like residues


## Detected protons



## Position of detected beam residues in coincidence with protons




# Preliminary results, detection of beam-like residues




Excitation energy




$5.7 < E^* < 6.1$  MeV




$7.0 < E^* < 7.4$  MeV



$7.8 < E^* < 8.2$  MeV




$8.6 < E^* < 9.0$  MeV







Sn

# Preliminary results for gamma and neutron-emission probabilities






# Origin of the structures



Calculations by Marc Dupuis

# Perspectives: measure simultaneously fission, neutron and gamma-emission probabilities




- Add fission detectors. **First time that fission is studied in a storage ring!**
- Demonstrate feasibility for measuring simultaneously  $P_f$ ,  $P_\gamma$  and  $P_n$ !
- Experiment accepted, to be probably conducted in 2024!
- After, produce **dedicated reaction chamber** to increase target-residue and fission detection efficiencies!

# Longer term perspectives: other stable & radioactive beams...

## Stable and radioactive beams at GSI/FAIR:

- $^{235}, ^{236}, ^{237}, ^{238}, ^{239}\text{U}$ ,  $^{235}, ^{236}, ^{237}, ^{238}\text{Pa}$ , etc
- Region around  $^{209}\text{Bi}$  and  $^{208}\text{Pb}$
- Region around  $^{136}\text{Xe}$  and  $^{135}\text{I}$
- Region of pre-actinides and actinides around shell  $N=126$




# Conclusions...

- Storage rings offer the ideal conditions to investigate surrogate reactions!
- First proof of principle experiment successfully conducted at the ESR in June 2022
  - $\Delta E^* \approx 600$  keV in accordance with expectations
  - Full separation and 70-100% detection efficiency for beam-like residues
  - Validation of new methodology for simultaneous measurement of  $P_\gamma$  and  $P_n$

## ...Perspectives

- Use  $P_\gamma$  and  $P_n$  to determine the neutron-induced cross sections of  $^{207}\text{Pb}$
- Add a fission detector to measure simultaneously  $P_\gamma$ ,  $P_n$  and  $P_f$  with  $^{238}\text{U}$  & target radius 0.5-1 mm.
- Build a dedicated reaction chamber to significantly increase efficiency for target residues and fission.
- Measurements with radioactive beams!

# Acknowledgements



This work is supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (ERC-Advanced grant NECTAR, grant agreement No 884715).

NECTAR: Nuclear rEaCTions At storage Rings



Prime 80 program from CNRS, PhD thesis of M. Sguazzin

**BACK-UP SLIDES**

# Angular uncertainty due to target radius

