



Electromagnetic moments in nuclei within nuclear DFT

Jacek Dobaczewski
University of York & University of Warsaw

Seminarium Fizyki Jądra Atomowego
Warszawa, 10 czerwca 2021



Jacek Dobaczewski

UNIVERSITY *of York*



UK Research
and Innovation



NucMagMom Collaboration (est. 2017)

- Michael Bender, Lyon
- Witek Nazarewicz, Mengzhi Chen, MSU
- Paolo Sassarini, Jérémy Bonard, York
- Ronald Fernando Garcia Ruiz, MIT

Literature

- B. Castel and I.S. Towner, *Modern theories of nuclear moments*, (Oxford Studies in Nuclear Physics) vol 12, ed P E Hodgson (Oxford: Clarendon,1990).
- Gerda Neyens, Rep. Prog. Phys. 66 (2003) 633–689.
- N.J. Stone, At. Data and Nucl. Data Tables 90 (2005) 75–176.
- M. Borrajo and J.L. Egido, Phys. Lett. B764 (2017) 328.
- L. Bonneau *et al.*, Phys. Rev. C91 (2015) 054307.
- O.I. Achakovskiy *et al.*, Eur. Phys. J. A (2014) 50:6.
- I. N. Borzov *et al.*, Phys. Atom. Nucl. 71 (2008) 469.



Jacek Dobaczewski

UNIVERSITY *of* York



UK Research
and Innovation



Outline

1. Recap on nuclear electromagnetic moments
2. Odd near doubly magic nuclei
3. N=83 isotones
4. Indium
5. Magnetic octupole moment in ^{45}Sc
6. Schiff moment in ^{225}Ra
7. ^{229}Th
8. Conclusions



Jacek Dobaczewski

UNIVERSITY *of* York



UK Research
and Innovation



Recap



Jacek Dobaczewski

UNIVERSITY *of York*



UK Research
and Innovation



Basic definitions

The electric and magnetic moments are defined as

$$Q_{\lambda\mu} = \langle \Psi | \hat{Q}_{\lambda\mu} | \Psi \rangle = \int q_{\lambda\mu}(\vec{r}) d^3\vec{r},$$

$$M_{\lambda\mu} = \langle \Psi | \hat{M}_{\lambda\mu} | \Psi \rangle = \int m_{\lambda\mu}(\vec{r}) d^3\vec{r},$$

where $|\Psi\rangle$ is a many-body state, and $q_{\lambda\mu}(\vec{r})$ and $m_{\lambda\mu}(\vec{r})$ are the corresponding electric and magnetic-moment densities:

$$q_{\lambda\mu}(\vec{r}) = e\rho(\vec{r})Q_{\lambda\mu}(\vec{r}),$$

$$m_{\lambda\mu}(\vec{r}) = \mu_N \left[g_s \vec{s}(\vec{r}) + \frac{2}{\lambda+1} g_l (\vec{r} \times \vec{j}(\vec{r})) \right] \cdot \vec{\nabla} Q_{\lambda\mu}(\vec{r}),$$

and e , g_s , and g_l are the elementary charge, and the spin and orbital gyromagnetic factors, respectively. The multipole functions (solid harmonics) have the standard form: $Q_{\lambda\mu}(\vec{r}) = r^\lambda Y_{\lambda\mu}(\theta, \phi)$.



Schmidt limits

The magnetic operator $\bar{\mu}$ is a one-body operator and the magnetic dipole moment μ is the expectation value of $\bar{\mu}_z$. The M1 operator acting on a composed state $|Im\rangle$ can then be written as the sum of single particle M1 operators $\bar{\mu}_z(j)$ acting each on an individual valence nucleon with total momentum j :

$$\mu = g_L \mathbf{L} + g_s \mathbf{s}$$

$$\mu(I) \equiv \left\langle I(j_1, j_2, \dots, j_n), m = I \left| \sum_{i=1}^n \bar{\mu}_z(i) \right| I(j_1, j_2, \dots, j_n), m = I \right\rangle \quad (2.1)$$

The single particle magnetic moment $\mu(j)$ for a valence nucleon around a doubly magic core is uniquely defined by the quantum numbers l and j of the occupied single particle orbit [22]:

$$\text{for an odd proton: } \left\{ \begin{array}{ll} \mu = j - \frac{1}{2} + \mu_p & \text{for } j = l + \frac{1}{2} \\ \mu = \frac{j}{j+1} \left(j + \frac{3}{2} - \mu_p \right) & \text{for } j = l - \frac{1}{2} \end{array} \right. \quad (2.2)$$

$$\text{for an odd neutron: } \left\{ \begin{array}{ll} \mu = \mu_n & \text{for } j = l + \frac{1}{2} \\ \mu = -\frac{j}{j+1} \mu_n & \text{for } j = l - \frac{1}{2} \end{array} \right. \quad (2.3)$$


**Schmidt
limits**

These single particle moments calculated using the free proton and free neutron moments ($\mu_p = +2.793$, $\mu_n = -1.913$) are called the Schmidt moments. In a nucleus, the magnetic



Experiment

M.G. Mayer and J.H.D. Jensen, *Elementary Theory of Nuclear Shell Structure*, (Wiley, New York, 1955)



Jacek Dobaczewski


UNIVERSITY of York



UK Research and Innovation



Experiment



M.G. Mayer and J.H.D. Jensen, Elementary Theory of Nuclear Shell Structure, (Wiley, New York, 1955)



Jacek Dobaczewski

UNIVERSITY of York



UK Research and Innovation



Odd near doubly magic nuclei



Jacek Dobaczewski


UNIVERSITY *of* York



UK Research
and Innovation



Odd near doubly magic nuclei



Jacek Dobaczewski

UNIVERSITY of York



UK Research and Innovation



Mechanism for e-m moments generation

● In nuclear DFT, properties of odd nuclei can be analysed in terms of the self-consistent polarisation effects caused by the presence of the unpaired nucleon.

● A non-zero quadrupole moment of the odd nucleon induces deformation of the total mean field and thus generates quadrupole moments of all remaining nucleons.

$$V = -\lambda Q_1 Q_2$$

● The latter moments enhance the deformation of the mean field even more, which in turn influences the quadrupole moment of the odd nucleon.

● In a self-consistent solution, these mutual polarisation are effectively summed up to infinity, whereupon the final total quadrupole deformation and electric quadrupole moment Q of the system are generated.

● A non-zero spin and current distributions of the odd particle influence those of all other nucleons and in the self-consistent solution lead to a specific polarisation of the system and its non-zero magnetic dipole moment μ .

$$V = -\lambda \sigma_1 \sigma_2$$

● All nucleons contribute to the moments Q and μ of the system, with individual contributions of nucleons depending on their individual polarisation responses to the deformed and polarised mean field.



Electric quadrupole moments



Jacek Dobaczewski

UNIVERSITY *of* York




UK Research
and Innovation



Electric quadrupole moments Q

P. Sassarini, J.D., J. Bonard, R.F. Garcia Ruiz, to be published



- Spectroscopic moments
- Average values for UNEDF1
SLy4
SkO'
D1S
N3LO
- Relative RMS deviations much smaller than the residuals



Jacek Dobaczewski

UNIVERSITY of York



UK Research and Innovation



Magnetic dipole moments



Jacek Dobaczewski

UNIVERSITY *of* York




UK Research
and Innovation



Magnetic dipole moments

P. Sassarini, J.D., J. Bonard, R.F. Garcia Ruiz, to be published



Jacek Dobaczewski

UNIVERSITY of York




UK Research and Innovation



Magnetic dipole moments vs. experiment

P. Sassarini, J.D., J. Bonard, R.F. Garcia Ruiz, to be published



Jacek Dobaczewski

UNIVERSITY of York




UK Research and Innovation



Optimisation of the spin-spin interaction

P. Sassarini, J.D., J. Bonard, R.F. Garcia Ruiz, to be published



Jacek Dobaczewski

UNIVERSITY of York




UK Research and Innovation



Magnetic dipole moments

P. Sassarini, J.D., J. Bonard, R.F. Garcia Ruiz, to be published



Jacek Dobaczewski

UNIVERSITY of York




UK Research and Innovation



Magnetic dipole moments

P. Sassarini, J.D., J. Bonard, R.F. Garcia Ruiz, to be published



Jacek Dobaczewski

UNIVERSITY of York



UK Research and Innovation



N=83 isotones





Jacek Dobaczewski

UNIVERSITY *of York*



UK Research
and Innovation





Indium



Jacek Dobaczewski

UNIVERSITY *of* York




UK Research
and Innovation



Magnetic dipole moments in indium

A.R. Vernon et al., submitted



Jacek Dobaczewski

UNIVERSITY of York




UK Research and Innovation



Electric quadrupole moments in indium

A.R. Vernon et al., submitted



Jacek Dobaczewski

UNIVERSITY of York



UK Research and Innovation



Magnetic octupole moment in ^{45}Sc



Jacek Dobaczewski

UNIVERSITY *of York*




UK Research
and Innovation




Visualisation of the magnetic multipole moments in axial symmetry


$\lambda=1$



$\lambda=2$



$\lambda=3$



Axial solid harmonics:

$\lambda\mu$	$Q_{\lambda\mu}$	$\nabla_z Q_{\lambda\mu}$
00	$\sqrt{\frac{1}{4\pi}}$	0
10	$\sqrt{\frac{3}{4\pi}}z$	$\sqrt{\frac{3}{4\pi}}$ = $\sqrt{3}Q_{00}$
20	$\sqrt{\frac{5}{16\pi}}(2z^2 - x^2 - y^2)$	$\sqrt{\frac{5}{\pi}}z$ = $\sqrt{\frac{20}{3}}Q_{10}$
30	$\sqrt{\frac{7}{16\pi}}(2z^3 - 3x^2z - 3y^2z)$	$\sqrt{\frac{7}{16\pi}}3(2z^2 - x^2 - y^2)$ = $\sqrt{\frac{63}{5}}Q_{20}$

Axial electric and magnetic-moment densities:

$$q_{\lambda 0}(r, \theta) = e\rho(r, \theta)Q_{\lambda 0}(r, \theta),$$

$$m_{\lambda 0}(r, \theta) = \mu_N \left[g_s s_z(r, \theta) + \frac{2}{\lambda+1} g_l (\vec{r} \times \vec{j})_z(r, \theta) \right] \cdot \nabla_z Q_{\lambda 0}(r, \theta),$$

or

$$m_{\lambda 0}(r, \theta) = \mu_N \left[g_s s_z(r, \theta) + \frac{2}{\lambda+1} g_l I_z(r, \theta) \right] C_\lambda Q_{(\lambda-1)0}(r, \theta),$$



Jacek Dobaczewski


UNIVERSITY of York



UK Research and Innovation



HF+AMP, magnetic moments in ^{45}Sc



R. P. de Groote *et al.*, arXiv:2005.00414



Schiff moment in ^{225}Ra



Jacek Dobaczewski


UNIVERSITY *of York*



UK Research
and Innovation



^{225}Ra Schiff moment vs. ^{225}Ra octupole moment



J.D., J. Engel, M. Kortelainen, P. Becker, *Phys. Rev. Lett.*, 121, 232501 (2018)



Jacek Dobaczewski


UNIVERSITY of York



UK Research and Innovation



^{225}Ra Schiff moment vs. ^{224}Ra octupole moment



J.D., J. Engel, M. Kortelainen, P. Becker, Phys. Rev. Lett., 121, 232501 (2018)



Jacek Dobaczewski

UNIVERSITY of York



UK Research and Innovation



^{229}Th



Jacek Dobaczewski

UNIVERSITY *of York*




UK Research
and Innovation



What is known about $^{229\text{m}}\text{Th}$?

Energies in keV



What we can aim for

1. Binding energies
2. **Odd-Even Mass Staggering**
3. **Axially and octupolly deformed**
4. Half-life
5. **Proton Quadrupole moments**
6. **Magnetic dipole moment**
7. Mean_square radii difference
8. **First few transition rates**

$$\Delta_n = 0.77 \text{ MeV}$$

$$\Delta_p = 0.68 \text{ MeV}$$

Octupole: degree of freedom

$$Q_{5/2} = 8.8(1) \text{ eb}$$


$$Q_{3/2} = 8.7(3) \text{ eb}$$

$$\mu_{5/2} = 0.360(7) \mu_N$$

$$\mu_{3/2} = -0.37(6) \mu_N$$



Reproduction of experimental odd-even mass staggering



Experimental values to reproduce:

$$\Delta_n = 0.77 \text{ MeV}$$

$$\Delta_p = 0.68 \text{ MeV}$$

Adjusted pairing

Interaction	$V_{0,n}$	$V_{0,p}$
SIII	181.15	220.19
SKM*	181.46	216.25
SKO'	163.82	184.34
SKXc	139.02	173.63
SLY4	207.76	231.89
UDF0	130.70	156.45
UDF1	145.35	169.80

Source:

<https://people.physics.anu.edu.au/~ecs103/chart/>

J.D., J. Bonnard, P. Becker, et al. to be published



Jacek Dobaczewski

UNIVERSITY of York



UK Research and Innovation





Nilsson levels in ^{229}Th vs. octupole deformation

X Let's go crossing hunting

SIII used before for ^{229}Th calculations:
E.Litvinova *et al.*, Phys. Rev.C, 79 064303 (2009)

Preliminary



Evolution of the energy of the blocked state with the octupole deformation in ^{229}Th

J.D., J. Bonnard, P. Becker, *et al.* to be published



Conclusions

1. Ground-state and isomeric electromagnetic moments are known in hundreds of odd and odd-odd nuclei, measured by atomic spectroscopic methods up to a **very high precision**.
2. In the standard shell-model calculations, agreement with data is achieved by using the concept of **effective charges and g-factors**.
3. In the nuclear DFT calculations, magnetic moments have been **rarely considered** so far.
4. Poorly known **time-odd sector** of the nuclear DFT crucially influences the magnetic moments.
5. **Adjustments of the nuclear DFT coupling constants to data** should take the magnetic moments into account.



Thank you



Jacek Dobaczewski

UNIVERSITY *of York*



UK Research
and Innovation



„Spin” magnetic dipole moment

In this study we use the single-particle magnetic-dipole-moment operator for neutron and proton bare orbital and spin gyrosopic factors,

$$g_\ell^p = \mu_N, \quad g_s^n = -3.826 \mu_N, \quad g_s^p = +5.586 \mu_N,$$

which reads

$$\hat{\mu} = g_\ell^p \hat{L}_p + g_s^n \hat{S}_n + g_s^p \hat{S}_p,$$

where \hat{L}_ν and \hat{S}_ν for $\nu = n, p$ are the operators of orbital and spin angular momenta, respectively. Since the total angular momentum $\hat{J} = \sum_{\nu=n,p} (\hat{L}_\nu + \hat{S}_\nu)$ is conserved, it is convenient to subtract its eigenvalue from the spectroscopic magnetic moments of odd- Z nuclei and to define "spin" magnetic moments μ^S as

$$\begin{aligned} \mu^S = \mu &= g_\ell^p \langle \hat{L}_p \rangle + g_s^n \langle \hat{S}_n \rangle + g_s^p \langle \hat{S}_p \rangle && \text{for } Z \text{ even,} \\ \mu^S = \mu - J \mu_N & && \\ &= g_\ell^{pn} \langle \hat{L}_n \rangle + g_s^{pn} \langle \hat{S}_n \rangle + g_s^{pp} \langle \hat{S}_p \rangle && \text{for } Z \text{ odd.} \end{aligned}$$


with

$$g_\ell^{pn} = -\mu_N, \quad g_s^{pn} = -4.826 \mu_N, \quad g_s^{pp} = +4.586 \mu_N.$$



HF+AMP, deformation energies in ^{45}Sc

R. P. de Groote et al., arXiv:2005.00414



isoscalar isovector Landau parameters g_0 & g_0'

$$V(\vec{r}_1, \vec{r}_2, \vec{r}_3, \vec{r}_4) = N_0 \left[g_0(\sigma_1 \cdot \sigma_2) + g_0'(\sigma_1 \cdot \sigma_2)(\tau_1 \cdot \tau_2) \right] \delta(\vec{r}_1 - \vec{r}_2)\delta(\vec{r}_1 - \vec{r}_3)\delta(\vec{r}_2 - \vec{r}_4)$$



HF + angular momentum projection (AMP)

The Hartree-Fock (HF) spin and current intrinsic densities read:

$$\vec{s}(\vec{r}) = \sum_{\sigma\sigma'} \vec{\sigma}_{\sigma'\sigma} \rho(\vec{r}\sigma, \vec{r}\sigma'), \quad \vec{j}(\vec{r}) = \frac{1}{2i} \sum_{\sigma} (\vec{\nabla} - \vec{\nabla}') \rho(\vec{r}\sigma, \vec{r}'\sigma),$$

where the one-body density matrix $\rho(\vec{r}\sigma, \vec{r}'\sigma')$ can be split into the core and odd-particle contributions:

$$\rho(\vec{r}\sigma, \vec{r}'\sigma') = \sum_{i=1}^{A-1} \psi_i(\vec{r}\sigma) \psi_i^*(\vec{r}'\sigma') + \psi_{\text{odd}}(\vec{r}\sigma) \psi_{\text{odd}}^*(\vec{r}'\sigma'),$$

and where $\psi(\vec{r}\sigma)$ are the self-consistent single-particle wave functions of occupied states. The HF wave function of an odd system $|\Phi\rangle = |\Phi^{\text{core}}\rangle \otimes |\psi^{\text{odd}}\rangle = \sum_I |\Psi_I\rangle$ has the conserved-angular-momentum components:

$$|\Psi_I\rangle = \sum_{J=0,2,4,\dots} \sum_{j=K,K+2,K+4,\dots} \left[|\Psi_J^{\text{core}}\rangle |\psi_j^{\text{odd}}\rangle \right]_I,$$

In ^{45}Sc , the angular-momentum projected ground state can be presented as:

$$\begin{aligned} |\Psi_{7/2}\rangle &= \left[|\Psi_0^{\text{core}}\rangle |\psi_{7/2}^{\text{odd}}\rangle \right]_{7/2} + \left[|\Psi_2^{\text{core}}\rangle |\psi_{7/2}^{\text{odd}}\rangle \right]_{7/2} \\ &+ \left[|\Psi_2^{\text{core}}\rangle |\psi_{11/2}^{\text{odd}}\rangle \right]_{7/2} + \left[|\Psi_4^{\text{core}}\rangle |\psi_{7/2}^{\text{odd}}\rangle \right]_{7/2} + \dots \end{aligned}$$

The first term represents a spherical core coupled to the spherical $j = 7/2$ wave function of the odd particle. The second term represents the lowest-order coupling of the odd-particle to the lowest $J = 2$ state of the core.



Who?

Jacek Dobaczewski, University of York, UK

Karim Bennaceur, Université Claude Bernard Lyon 1, F

Ruben de Groote, University of Jyväskylä, FI

Markus Kortelainen, University of Jyväskylä, FI

Ronald Fernando Garcia Ruiz, MIT, Cambridge, USA

Iain Moore, University of Jyväskylä, FI

Gerda Neyens, KU Leuven and CERN, B and CH

Alessandro Pastore, University of York, UK

What?

We propose **supporting European facilities** involved in precision **measurements of nuclear moments** with novel advanced modelling of these observables within nuclear **density-functional-theory (DFT) approaches**.

ADVANCING FRONTIER KNOWLEDGE



Jacek Dobaczewski

UNIVERSITY *of York*



UK Research
and Innovation

